

CUP-LENGTH ESTIMATES FOR LEAF-WISE INTERSECTIONS

PETER ALBERS AND AL MOMIN

ABSTRACT. We prove that on a restricted contact type hypersurface the number of leaf-wise intersections is bounded from below by a certain cup-length.

1. INTRODUCTION

Let (M, ω) be a symplectic manifold and $\Sigma \subset M$ be a hypersurface. Then Σ is foliated by the characteristic foliation induced by the line bundle $\ker \omega|_{\Sigma} \rightarrow \Sigma$. We denote by L_x the leaf through $x \in \Sigma$. Let $\psi \in \text{Ham}(M)$ be a Hamiltonian diffeomorphism. Then a leaf-wise intersection is a point $x \in \Sigma$ with the property that $\psi(x) \in L_x$.

Definition 1.1. Let $\iota : \Sigma \hookrightarrow M$ be the inclusion map.

(1) We denote by

$$\text{cl}(\Sigma, M) := \max\{k \mid \exists a_1, \dots, a_k \in H^{\geq 1}(M; \mathbb{Z}/2) \text{ with } \iota^*(a_1 \cup \dots \cup a_k) \neq 0\} \quad (1.1)$$

the relative cup-length of Σ in M .

- (2) Suppose $\omega = d\lambda$ and that Σ is of restricted contact type, i.e. $\alpha := \iota^*\lambda$ is a contact form on Σ . Then we denote by $\wp(\Sigma, \alpha) > 0$ the minimal period of a Reeb orbit of (Σ, α) which is contractible in M . If there exists no such Reeb orbit we set $\wp(\Sigma, \alpha) = \infty$.
- (3) We denote by $\text{Ham}_c(M)$ the space of Hamiltonian diffeomorphisms generated by compactly supported, time dependent Hamiltonian functions and by $\|\psi\|$ the Hofer norm.
- (4) (M, ω) is called convex at infinity if it is isomorphic to the symplectization of a compact contact manifold at infinity.

Theorem 1. *Let $\Sigma \subset (M, d\lambda)$ be a closed, bounding, restricted contact type hypersurface and $(M, d\lambda)$ be convex at infinity. If $\psi \in \text{Ham}_c(M)$ satisfies $\|\psi\| < \wp(\Sigma, \alpha)$ then*

$$\nu_{\text{leaf}}(\psi) := \#\{\text{leaf-wise intersections of } \psi\} \geq \text{cl}(\Sigma, M) + 1. \quad (1.2)$$

Remark 1.2. The search for leaf-wise intersections was initiated by Moser in [Mos78] and pursued further in [Ban80, Hof90, EH89, Gin07, Dra08, AF10a, Zil08, AF08, AM09, Gür09, Kan09, AF10b, AF10c, Mer10]. We refer to [AF08] for a brief history of the problem.

Remark 1.3. The example $S^1 \subset \mathbb{C}$ shows that if $\|\psi\| > \wp(\Sigma, \alpha)$ then $\nu_{\text{leaf}}(\psi) = 0$ is possible.

Remark 1.4. The relative cup-length is smaller than the cup-length of M and Σ . In the case $\Sigma = S^*B \subset T^*B$ is a unit cotangent bundle we have $\text{cl}(S^*B, T^*B) \geq \text{cl}(B) - 1$ by examining the Gysin sequence.

Remark 1.5. For simplicity we shall use $\mathbb{Z}/2$ -coefficients for all (co-)homology theories in this paper. We expect that Theorem 1 continues to hold with $\mathbb{Z}$ -coefficients.

2000 *Mathematics Subject Classification.* 53D40, 37J10, 58f05.

Key words and phrases. Rabinowitz Floer homology, leaf-wise intersections, cup-length estimates.

Remark 1.6. Cup-length estimates have been established for Lagrangian intersections by Floer [Flo89], Hofer [Hof88], and Liu [Liu05] in terms of the cup-length of the Lagrangian submanifold and for fixed points of Hamiltonian diffeomorphisms by Schwarz [Sch98] in terms of the quantum cup-length of the symplectic manifold.

2. LEAF-WISE INTERSECTIONS AND THE RABINOWITZ ACTION FUNCTIONAL

Let (M, ω) be a symplectic manifold and $f \in C^\infty(M)$ an autonomous Hamiltonian function. Since energy is preserved the hypersurface $\Sigma := f^{-1}(0)$ is invariant under the Hamiltonian flow ϕ_f^t of f . The Hamiltonian flow ϕ_f^t is generated by the Hamiltonian vector field X_f which is uniquely defined by the equation $\omega(X_f, \cdot) = df$. If 0 is a regular value of f the hypersurface is a coisotropic submanifold which is foliated by 1-dimensional isotropic leaves, see [MS98, Section 3.3]. If we denote by L_x the leaf through $x \in \Sigma$ we have the equality

$$L_x = \bigcup_{t \in \mathbb{R}} \phi_f^t(x). \quad (2.1)$$

Given a time-dependent Hamiltonian function $H : [0, 1] \times M \rightarrow \mathbb{R}$ with Hamiltonian flow ϕ_H^t we are interested in points $x \in \Sigma$ with the property

$$\phi_H^1(x) \in L_x. \quad (2.2)$$

This notion was introduced and studied by Moser in [Mos78]. Such points are called leaf-wise intersections. We recall some notions from [AF10a].

Definition 2.1. A leaf-wise intersection $x \in \Sigma$ is called periodic if the leaf L_x is a closed orbit of the flow ϕ_f^t .

Definition 2.2. A pair $\mathfrak{M} = (F, H)$ of Hamiltonian functions $F, H : S^1 \times M \rightarrow \mathbb{R}$ is called a Moser pair if it satisfies

$$F(t, \cdot) = 0 \quad \forall t \in [0, \tfrac{1}{2}] \quad \text{and} \quad H(t, \cdot) = 0 \quad \forall t \in [\tfrac{1}{2}, 1], \quad (2.3)$$

and F is of the form $F(t, x) = \rho(t)f(x)$ for some smooth map $\rho : S^1 \rightarrow [0, \infty)$ with $\int_0^1 \rho(t)dt = 1$ and $f : M \rightarrow \mathbb{R}$.

Definition 2.3. We set

$$\mathcal{H} := \{H \in C^\infty(S^1 \times M) \mid H \text{ has compact support and } H(t, \cdot) = 0 \quad \forall t \in [\tfrac{1}{2}, 1]\} \quad (2.4)$$

Remark 2.4. It is easy to see that the $\text{Ham}(M, \omega) \equiv \{\phi_H^1 \mid H \in \mathcal{H}\}$, e.g. [AF10a].

Let $(M, \omega = d\lambda)$ be an exact symplectic manifold. Then for a Moser pair $\mathfrak{M} = (F, H)$ the perturbed Rabinowitz action functional is defined by

$$\begin{aligned} \mathcal{A}^{\mathfrak{M}} : \mathcal{L}_M \times \mathbb{R} &\longrightarrow \mathbb{R} \\ (v, \eta) &\mapsto - \int_{S^1} v^* \lambda - \int_0^1 H(t, v) dt - \eta \int_0^1 F(t, v) dt \end{aligned} \quad (2.5)$$

where $\mathcal{L}_M := C^\infty(S^1, M)$. A critical point (v, η) of $\mathcal{A}^{\mathfrak{M}}$ is a solution of

$$\left. \begin{aligned} \partial_t v &= \eta X_F(t, v) + X_H(t, v) \\ \int_0^1 F(t, v) dt &= 0 \end{aligned} \right\} \quad (2.6)$$

In [AF10a] it is proved that critical points of $\mathcal{A}^{\mathfrak{M}}$ give rise to leaf-wise intersections.

Proposition 2.5 ([AF10a]). *Let (v, η) be a critical point of $\mathcal{A}^{\mathfrak{M}}$. Then $x := v(0) \in f^{-1}(0)$ and*

$$\phi_H^1(x) \in L_x \quad (2.7)$$

thus, x is a leaf-wise intersection.

Moreover, the map $\text{Crit}\mathcal{A}^{\mathfrak{M}} \rightarrow \{\text{leaf-wise intersections}\}$ is injective unless there exists a periodic leaf-wise intersection (see Definition 2.1).

Definition 2.6. A Moser pair $\mathfrak{M} = (F, H)$ is of restricted contact type if the following four conditions hold.

- (1) 0 is a regular value of f .
- (2) df has compact support.
- (3) The hypersurface $f^{-1}(0)$ is a closed restricted contact type hypersurface of (M, λ) .
- (4) The Hamiltonian vector field X_f restricts to the Reeb vector field on $f^{-1}(0)$.

Remark 2.7. If $\Sigma \subset (M, d\lambda)$ is a closed, bounding, restricted contact type hypersurface then it is easy to construct a restricted contact type Moser pair $\mathfrak{M}_0 = (F, 0)$ with $\Sigma = f^{-1}(0)$. We fix such a Moser pair for the rest of the paper. Critical points (v, η) of the unperturbed Rabinowitz action functional $\mathcal{A}^{(F,0)}$ are η -periodic Reeb orbits in Σ or if $\eta = 0$ constant loops in Σ . Their critical values are $\mathcal{A}^{(F,0)}(v, \eta) = -\eta$.

In [CF09] Cieliebak and Frauenfelder construct a Floer homology for the unperturbed Rabinowitz action functional. This has been extended to the perturbed case by the first author and Frauenfelder in [AF10a]. The corresponding Floer equation for maps $u : \mathbb{R} \times S^1 \rightarrow M$ and $\eta : \mathbb{R} \rightarrow \mathbb{R}$ is

$$\left. \begin{aligned} \partial_s u + J(s, t, u)(\partial_t u - \eta X_F(t, u) - X_{H_s}(t, u)) &= 0 \\ \partial_s \eta - \int_0^1 F(t, u) dt &= 0. \end{aligned} \right\} \quad (2.8)$$

Here J is a smooth (s, t) -dependent family of compatible almost complex structures and $H_s : S^1 \times M \rightarrow \mathbb{R}$ is a smooth s -dependent family of functions. Counting solutions of the s -independent equation modulo $\mathbb{R}$ -action defines the boundary operator in Rabinowitz Floer homology. In this paper we do not need the full machinery of Rabinowitz Floer homology.

We set $w = (u, \eta)$ for solutions of (2.6). We will think of w also as a map $w : \mathbb{R} \rightarrow \mathcal{L}_M \times \mathbb{R}$.

Definition 2.8. The energy of a map $w = (u, \eta)$ is defined as

$$E(w) := \int_{-\infty}^{\infty} \int_0^1 \|\partial_s u\|^2 dt ds + \int_{-\infty}^{\infty} |\partial_s \eta|^2 ds. \quad (2.9)$$

The following has been established in [AF10a].

Lemma 2.9 ([AF10a], Lemma 2.7). Let w solve (2.8) with limits

$$w(\pm\infty) := \lim_{s \rightarrow \pm\infty} w(s) \in \text{Crit}\mathcal{A}^{(F, H_{\pm})} \quad (2.10)$$

then we have

$$E(w) \leq \mathcal{A}^{(F, H_-)}(w(-\infty)) - \mathcal{A}^{(F, H_+)}(w(+\infty)) - \int_{-\infty}^{\infty} \int_0^1 \min_M \partial_s H_s(t, \cdot) dt ds \quad (2.11)$$

and equality holds if $\partial_s H_s = 0$.

Theorem 2.10. *Let $w_n = (u_n, \eta_n)$ be a sequence of solutions of (2.8) for which there exists $a < b$ such that*

$$a \leq \mathcal{A}^{(F,H)}(w_n(s)) \leq b \quad \forall s \in \mathbb{R}. \quad (2.12)$$

Then for every reparametrisation sequence $(\sigma_n) \subset \mathbb{R}$ the sequence $u_n(\cdot + \sigma_n)$ has a subsequence which converges in $C_{\text{loc}}^\infty(\mathbb{R} \times S^1, M)$ and similarly for $\eta_n(\cdot + \sigma_n)$.

3. MODULI SPACES

Let $k \geq 1$ be a natural number. We choose a smooth family of functions $\beta_R \in C^\infty(\mathbb{R}, [0, 1])$ satisfying

- (1) for $R \geq 1$: $\beta_R'(s) \cdot s \leq 0$ for all $s \in \mathbb{R}$, $\beta_R(s) = 1$ for $s \in [0, (k+1)R]$, and $\beta_R(s) = 0$ for $s \leq -1$ and $s \geq (k+1)R + 1$,
- (2) for $R \leq 1$: $\beta_R(s) \leq R$ for all $s \in \mathbb{R}$ and $\text{supp} \beta_R \subset [-1, k+2]$,
- (3) $\lim_{R \rightarrow \infty} \beta_R(s) =: \beta_\infty^+(s)$ and $\lim_{R \rightarrow \infty} \beta_R(s + (k+1)R) =: \beta_\infty^-(s)$ exists, where the limit is taken with respect to the C_{loc}^∞ topology.




FIGURE 1. The function β_R .

We fix a Hamiltonian function $H : [0, 1] \times M \rightarrow \mathbb{R}$ with $H(t, \cdot) = 0$ for all $t \in [\frac{1}{2}, 1]$ and set $K_R(s, t, x) := \beta_R(s)H(t, x)$ and

$$\mathfrak{M}_R := (F, K_R(s, t, x)). \quad (3.1)$$

For every $R \geq 0$ we define

$$\mathcal{M}(R) := \left\{ w = (u, \eta) \in C^\infty(\mathbb{R}, \mathcal{L}_M \times \mathbb{R}) \left| \begin{array}{l} w \text{ solves (2.8) with } \mathfrak{M}_R \\ \lim_{s \rightarrow \pm\infty} u(s) \in \Sigma \end{array} \right. \right\}. \quad (3.2)$$

and

$$\mathcal{M}[0, R] := \{(r, w) \mid 0 \leq r \leq R \text{ and } w \in \mathcal{M}(r)\}. \quad (3.3)$$

see figure 2.

Lemma 3.1. For $w \in \mathcal{M}(R)$ we have

$$E(w) \leq \|H\|. \quad (3.4)$$

Moreover, for $R = 0$ we have $E(w) = 0$. Furthermore,

$$-2 \int_0^1 \max_M |H(t, \cdot)| dt \leq \mathcal{A}^{\mathfrak{M}_R(s)}(w(s)) \leq 2 \int_0^1 \max_M |H(t, \cdot)| dt \quad (3.5)$$

holds.

PROOF. We compute using $\partial_s w + \nabla \mathcal{A}^{\mathfrak{M}_R(s)} = 0$

$$\begin{aligned}
\underbrace{\mathcal{A}^{\mathfrak{M}_0}(w(-\infty))}_{=0} - \mathcal{A}^{\mathfrak{M}_R(s)}(w(s)) &= - \int_{-\infty}^s \frac{d}{ds} \mathcal{A}^{\mathfrak{M}_R(s)}(w(s)) ds \\
&= - \int_{-\infty}^s d\mathcal{A}^{\mathfrak{M}_R(s)}(w(s)) \cdot \partial_s w ds - \int_{-\infty}^s \frac{\partial \mathcal{A}^{\mathfrak{M}_R(s)}}{\partial s}(w(s)) ds \\
&= - \int_{-\infty}^s \underbrace{(\nabla \mathcal{A}^{\mathfrak{M}_R(s)})}_{=-\partial_s w} \cdot \partial_s w ds + \int_{-\infty}^s \int_0^1 \frac{\partial K_R}{\partial s}(s, t, u(s, t)) dt ds \\
&= \int_{-\infty}^s \|\partial_s w(s)\|^2 ds + \int_{-\infty}^s \int_0^1 \frac{\partial K_R}{\partial s}(s, t, u(s, t)) dt ds \\
&\geq \int_{-\infty}^s \left(\frac{\partial}{\partial s} \beta_R(s) \right) \int_0^1 H(t, u(s, t)) dt ds \\
&\geq \underbrace{\int_{-\infty}^0 \left(\frac{\partial}{\partial s} \beta_R(s) \right) ds}_{=1} \cdot \int_0^1 -\max_M |H(t, \cdot)| dt \\
&\quad + \underbrace{\int_0^\infty \left(\frac{\partial}{\partial s} \beta_R(s) \right) ds}_{=-1} \cdot \int_0^1 \max_M |H(t, \cdot)| dt \\
&\geq -2 \int_0^1 \max_M |H(t, \cdot)| dt
\end{aligned}$$

The other inequality is proved by replacing $\int_{-\infty}^s$ by $\int_s^\infty$. The first two assertions follow from




FIGURE 2. An element of the moduli space $\mathcal{M}(R)$.

Lemma 2.9 together with the observation that $\mathcal{A}^{(F, \beta_R H)}(w(\pm\infty)) = 0$ and

$$\begin{aligned}
-\int_{-\infty}^{\infty} \int_0^1 \min_M \partial_s K_R(s, t, \cdot) dt ds &= -\int_{-\infty}^{\infty} \int_0^1 \min_M \{\partial_s \beta_R(s) H(t, \cdot)\} dt ds \\
&= -\int_{-\infty}^0 \int_0^1 \partial_s \beta_R(s) \min_M H(t, \cdot) dt ds \\
&\quad - \int_0^{+\infty} \int_0^1 \partial_s \beta_R(s) \max_M H(t, \cdot) dt ds \\
&= \beta_R(0) \|H\| \leq \|H\|.
\end{aligned} \tag{3.6}$$

□

Proposition 3.2. *For any $R \geq 0$ the moduli spaces $\mathcal{M}(R)$ and $\mathcal{M}[0, R]$ are compact. Moreover, $\mathcal{M}(0) \cong \Sigma$.*

PROOF. This follows easily as in the proof of Theorem A in [AF10a] as follows. Using Lemma 3.1 we can apply Theorem 2.10 to extract C_{loc}^∞ -convergent subsequences of any sequence $w_n(s - \sigma_n)$ where $w_n \in \mathcal{M}(R)$ and $(\sigma_n) \subset \mathbb{R}$. Then it is proved in [AF10a] that if the sequence does not converge to an element in $\mathcal{M}(R)$ there has to exist a non-constant gradient flow line v of $\mathcal{A}^{(F, 0)}$ with one asymptotic end on Σ . Therefore, there exists a Reeb orbit x of period η on (Σ, α) which is contractible in M . We conclude

$$E(v) \leq \limsup E(w_n) \leq \|H\| < \wp(\Sigma, \alpha). \tag{3.7}$$

On the other hand we can compute $E(v)$ by Lemma 2.9:

$$E(v) = |\eta| \geq \wp(\Sigma, \alpha) \tag{3.8}$$

where the inequality follows from the definition of $\wp(\Sigma, \alpha)$. This contradiction shows that $\mathcal{M}(R)$ is compact. That $\mathcal{M}[0, R]$ is compact follows in the same way.

If $R = 0$ then according to Lemma 3.1 $E(w) = 0$ for all $w \in \mathcal{M}(0)$. So $\partial_s w(s) = 0$ and $w(s) = (p, 0) \in \text{Crit} \mathcal{A}^{(F, 0)}$ with $p \in \Sigma$ being the constant loop; thus, $\mathcal{M}(0) \cong \Sigma$. □

Remark 3.3. The moduli space $\mathcal{M}(R)$ is the zero-set of a Fredholm section $\mathcal{F}(R)$ of a Banach space bundle. The Fredholm index of $\mathcal{F}(R)$ equals $\text{ind } \mathcal{F}(R) = \dim \Sigma$. The moduli space $\mathcal{M}(0)$ contains only constant gradient flow lines with Lagrange multiplier $\eta = 0$, i.e. $\mathcal{M}(0) \cong \Sigma$. As shown in [AF10a] the Morse-Bott property of the defining function F implies that the Fredholm section is transverse to the zero-section for $R = 0$. In particular, $\mathcal{M}(0)$ is a smooth manifold. Since $\mathcal{M}(R)$ is compact it is a smooth manifold for R sufficiently small since transversality is an open property.

4. COHOMOLOGY OPERATIONS

We fix a natural number $k \geq 1$. The moduli space $\mathcal{M}(R)$ carries an evaluation map

$$\begin{aligned}
\text{ev}_R : \mathcal{M}(R) &\longrightarrow M^k = M \times \dots \times M \\
w = (u, \eta) &\mapsto (u(R, 0), u(2R, 0), \dots, u(kR, 0)).
\end{aligned} \tag{4.1}$$

For $R = 0$ the evaluation map is under the identification $\mathcal{M}(0) \cong \Sigma$ the diagonal embedding $\Sigma \hookrightarrow \Delta_\Sigma \subset M^k$. Similarly, $\mathcal{M}[0, R]$ has an evaluation map $\text{EV}(r, w) := \text{ev}_r(w)$ to M^k .

Next, we define a Morse theoretic realization of the cohomology operation

$$\begin{aligned} \Theta : H^*(M) \otimes \dots \otimes H^*(M) \otimes H_*(\Sigma) &\longrightarrow H_*(\Sigma) \\ a_1 \otimes \dots \otimes a_k \otimes b &\mapsto \iota^*(a_1 \cup \dots \cup a_k) \cap b \end{aligned} \quad (4.2)$$

where $\iota : \Sigma \hookrightarrow M$ is the inclusion map.

Remark 4.1. Since the symplectic manifold (M, ω) is exact it either has boundary or is non-compact. In the following we will choose Morse functions on M in order to model singular (co-)homology by Morse (co-)homology. For this we need to restrict to a certain class of Morse functions. We are always considering the negative gradient flow. If M has boundary then we assume that the (positive) gradient of the Morse function points outward along the boundary of M . If M is non-compact we assume that the Morse function is proper and bounded from below. Under these assumptions standard Morse (co-)homology can be defined and is isomorphic to singular (co-)homology. From now on we assume whenever we choose Morse functions on M they are in the just prescribed class.

For that we choose Morse functions $f_1, \dots, f_k : M \rightarrow \mathbb{R}$ and $f_* : \Sigma \rightarrow \mathbb{R}$ and Riemannian metrics $g_1, \dots, g_k, g_*$. We set for critical points $x_j \in \text{Crit}(f_j)$ and $x_*^\pm \in \text{Crit}(f_*)$

$$\mathcal{M}(0, x_1, \dots, x_k, x_*^-, x_*^+) := \left\{ w = (u, \eta) \in \mathcal{M}(0) \left| \begin{array}{l} u(-\infty) \in W^u(x_*, f_*), u(+\infty) \in W^s(x_*, f_*) \\ \text{ev}_0(u) \in W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k) \end{array} \right. \right\}$$

see figure 3. In particular, $\mathcal{M}(0, x_1, \dots, x_k, x_*^\pm) = \emptyset$ unless $\bigcap W^s(x_i, f_i) \cap \Sigma \neq \emptyset$. We denote by $\text{CM}^*(f)$ the Morse chain complex associated to a Morse-Smale pair (f, g) .




FIGURE 3. An element of the moduli space $\mathcal{M}(0, x_1, \dots, x_k, x_*^-, x_*^+)$.

Proposition 4.2. *For generic Morse functions f_i and Riemannian metrics g_i the moduli space $\mathcal{M}(0, x_1, \dots, x_k, x_*^-, x_*^+)$ is a smooth manifold. Moreover, the map defined by*

$$\begin{aligned} \theta_0 : \text{CM}^*(f_1) \otimes \dots \otimes \text{CM}^*(f_k) \otimes \text{CM}_*(f_*) &\longrightarrow \text{CM}_*(f_*) \\ x_1 \otimes \dots \otimes x_k \otimes x_*^- &\mapsto \sum_{x_*^+} \#_2 \mathcal{M}(0, x_1, \dots, x_k, x_*^-, x_*^+) \cdot x_*^+ \end{aligned} \quad (4.3)$$

defines a chain map which on homology agrees with the cohomology operation Θ .

Remark 4.3. $\#_2 \mathcal{M}$ denotes the parity of the set $\mathcal{M}$ if it is finite and zero otherwise.

PROOF. It follows from standard Morse theory that for generic Morse functions f_i and Riemannian metrics g_i we have

$$W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k) \cap \Delta_\Sigma \subset M^k \quad (4.4)$$

where $\Delta_\Sigma \subset M^k$ is the diagonal embedding of Σ . Since for $R = 0$ the evaluation map $\text{ev}_0 : \mathcal{M}(0) \rightarrow M^k$ is this diagonal embedding we conclude

$$W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k) \cap \text{ev}_0. \quad (4.5)$$

Finally, choosing f_* and g_* generic we see that the moduli space $\mathcal{M}(0, x_1, \dots, x_k, x_*^-, x_*^+)$ is smooth. That $\theta_0 = \Theta$ after identifying Morse homology with singular homology is again standard Morse theory, see [Sch93]. $\square$

Since the moduli space $\mathcal{M}(R)$ is the zero-set of a Fredholm section $\mathcal{F}(R)$ and since $\mathcal{M}(R)$ is compact we can choose an arbitrarily small abstract perturbation of the Fredholm section $\mathcal{F}(R)$ such that the zero-set $\widetilde{\mathcal{M}}(R)$ of the perturbed Fredholm section $\widetilde{\mathcal{F}}(R)$ is a smooth compact finite-dimensional manifold. Since $\mathcal{M}(R)$ is already transverse for sufficiently small R we can arrange that $\widetilde{\mathcal{M}}(R) = \mathcal{M}(R)$ for small R . We point out that $\mathcal{M}[0, R]$ has a natural projection to $[0, R]$. The same abstract perturbation procedure gives rise to smooth perturbed moduli spaces $\widetilde{\mathcal{M}}[0, R]$ where the perturbation can be chosen with fixed ends, i.e., the fibers over 0 resp. R are $\mathcal{M}(0)$ resp. $\widetilde{\mathcal{M}}(R)$. The perturbed moduli space $\widetilde{\mathcal{M}}(R)$ resp. $\widetilde{\mathcal{M}}[0, R]$ still carries an evaluation map $\widetilde{\text{ev}}(R) : \widetilde{\mathcal{M}}(R) \rightarrow M^k$ resp. $\widetilde{\text{EV}} : \widetilde{\mathcal{M}}[0, R] \rightarrow M^k$.

For Morse functions $f_1, \dots, f_k : M \rightarrow \mathbb{R}$ and $f_* : \Sigma \rightarrow \mathbb{R}$, Riemannian metrics $g_1, \dots, g_k, g_*$, and critical points $x_j \in \text{Crit}(f_j)$ and $x_*^\pm \in \text{Crit}(f_*)$ we set

$$\widetilde{\mathcal{M}}(R, x_1, \dots, x_k, x_*^-, x_*^+) := \left\{ w = (u, \eta) \in \widetilde{\mathcal{M}}(R) \mid \begin{array}{l} u(-\infty) \in W^u(x_*^-, f_*), u(+\infty) \in W^s(x_*^+, f_*) \\ \widetilde{\text{ev}}_R(u) \in W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k) \end{array} \right\}$$

see figure 4.




FIGURE 4. An element of the moduli space $\widetilde{\mathcal{M}}(R, x_1, \dots, x_k, x_*^-, x_*^+)$.

Proposition 4.4. *For generic Morse functions f_i and Riemannian metrics g_i the moduli space $\widetilde{\mathcal{M}}(R, x_1, \dots, x_k, x_*^-, x_*^+)$ is a smooth manifold. Moreover, the map defined by*

$$\begin{aligned} \theta_R : \text{CM}^*(f_1) \otimes \dots \otimes \text{CM}^*(f_k) \otimes \text{CM}_*(f_*) &\longrightarrow \text{CM}_*(f_*) \\ x_1 \otimes \dots \otimes x_k \otimes x_*^- &\mapsto \sum_{x_*^+} \#_2 \widetilde{\mathcal{M}}(R, x_1, \dots, x_k, x_*^-, x_*^+) \cdot x_*^+ \end{aligned} \quad (4.6)$$

defines a chain map which is chain homotopic to the cohomology operation θ_0 .

PROOF. For generic Morse functions f_i , generic Riemannian metrics g_i , and generic perturbation of the Fredholm section we have

$$W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k) \pitchfork \widetilde{\text{ev}}_R. \quad (4.7)$$

For generic f_* and g_* the stable and unstable manifolds are transversal to the evaluation maps $u \mapsto u(\pm\infty)$ and thus the moduli space $\widetilde{\mathcal{M}}(R, x_1, \dots, x_k, x_*^-, x_*^+)$ is smooth. The map θ_R is a chain map by standard Morse theory since $\widetilde{\mathcal{M}}(R)$ is compact.

To prove that θ_R is chain homotopic to θ_0 we recall that $\widetilde{\mathcal{M}}[0, R]$ is perturbed while keeping the ends $\mathcal{M}(0)$ and $\widetilde{\mathcal{M}}(R)$ fixed. Perturbing further (with fixed ends), the evaluation map $\widetilde{\text{EV}} : \widetilde{\mathcal{M}}[0, R] \rightarrow M^k$ will be transverse to all products of unstable manifolds. Thus, the moduli space $\widetilde{\mathcal{M}}[0, R]$ together with $\widetilde{\text{EV}}$ induces a cobordism between the moduli spaces $\widetilde{\mathcal{M}}(R, x_1, \dots, x_k, x_*^\pm)$ and $\widetilde{\mathcal{M}}(0, x_1, \dots, x_k, x_*^\pm)$. This give rise to a chain homotopy operator between θ_0 and θ_R . $\square$

5. PROOF OF THEOREM 1

We assume that ψ has only finitely many leaf-wise intersections since otherwise we are done. We set $k := \text{cl}(\Sigma, M)$. It has been proved in [AF10a] that $\nu_{\text{leaf}}(\psi) \geq 1$ if $\|\psi\| < \wp(\Sigma, \alpha)$. Thus, we may assume that $k \geq 1$. We choose a Hamiltonian $H : [0, 1] \times M \rightarrow \mathbb{R}$ with $H(t, \cdot) = 0$ for $t \in [\frac{1}{2}, 1]$ such that $\phi_H^1 = \psi$. Let $\mathcal{LI} \subset \Sigma$ be the set of leaf-wise intersections of ψ .

We choose Morse functions $f_1, \dots, f_k$ and Riemannian metrics $g_1, \dots, g_k$ on M and f_* , g_* on Σ with the following properties

- (1) For $x_i \in \text{Crit}(f_i)$ with Morse index $\mu_{\text{Morse}}(x_i, f_i) \neq 0$ we have $W^s(x_i, f_i) \cap \mathcal{LI} = \emptyset$.
- (2) For all $n \in \mathbb{N}$ the evaluation maps $\widetilde{\text{ev}}_n : \widetilde{\mathcal{M}}(n) \rightarrow M^k$ are transverse to the products of stable manifolds of the (f_i, g_i) .
- (3) For all $n \in \mathbb{N}$ the evaluation maps at $\pm\infty$ are transverse to all stable and unstable manifolds of (f_*, g_*) .

The second and third property holds for a fixed n for a Baire set of Morse functions and Riemannian metrics as explained in the proofs of Propositions 4.2 and 4.4. Intersecting these Baire sets over $n \in \mathbb{N}$ we see (2) and (3) is a generic property. Avoiding finitely many unstable manifolds not of top dimension is clearly also a generic condition.

By definition of k we find cohomology classes $a_1, \dots, a_k \in H^{\geq 1}(M)$ with $\iota^*(a_1 \cup \dots \cup a_k) \neq 0 \in H^*(\Sigma)$. Thus, the cohomology operation Θ , see equation (4.2), is non-zero. According to Propositions 4.2 and 4.4 the cohomology operations θ_R then have to be non-zero for all $R \geq 0$. Thus, we can find critical points $x_i \in \text{Crit}(f_i)$ with Morse index $\mu_{\text{Morse}}(x_i, f_i) \neq 0$ and $x_*^\pm \in \text{Crit}(f_*)$ such that for all $n \in \mathbb{N}$ the moduli spaces

$$\widetilde{\mathcal{M}}(n, x_1, \dots, x_k, x_*^-, x_*^+) \neq \emptyset. \quad (5.1)$$

In fact, the choice of $x_i \in \text{Crit}(f_i)$ will depend on n in general. For notational convenience we suppress the n . In fact, there exists a subsequence n_k for which we can choose x_i fixed. We conclude

$$\widetilde{\mathcal{M}}(n) \neq \emptyset. \quad (5.2)$$

If the unperturbed moduli space $\mathcal{M}(n)$, see equation (3.2), were empty then for sufficiently small perturbations of the Fredholm section also the moduli space $\widetilde{\mathcal{M}}(n)$ would be empty. Indeed, if $\mathcal{M}(n) = \emptyset$ then the corresponding Fredholm section is transverse. Since $\mathcal{M}(n)$ is compact a sufficiently small perturbation of the Fredholm section remains empty and thus transverse. Therefore, we conclude that

$$\mathcal{M}(n) \neq \emptyset. \quad (5.3)$$

Now we can choose a sequence $w_n \in \mathcal{M}(n)$, that is, w_n solves equation (2.8) with Hamiltonian perturbation $\beta_n(s)H(t, x)$. Since $\widetilde{\mathcal{M}}(n) \cap_{\widetilde{\text{ev}}_n} (W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k)) \neq \emptyset$ and all evaluation maps are transverse a similar argument as above allows us to conclude that

$$\text{ev}_n(w_n) \in W^s(x_1, f_1) \times \dots \times W^s(x_k, f_k). \quad (5.4)$$

By Lemma 3.1 the action $\mathcal{A}^{\mathfrak{M}_n}(w_n(s))$ is uniformly bounded and we can apply Theorem 2.10. We consider the following sequences

$$w_n(s + jn), \quad j = 0, \dots, k+1. \quad (5.5)$$

For $j = 1, \dots, k$ these sequences converge (after choosing a subsequence) to solutions $\varpi^j = (u^j, \eta^j) : \mathbb{R} \rightarrow \mathcal{L}_M \times \mathbb{R}$ of

$$\left. \begin{aligned} \partial_s u^j + J(s, t, u^j) (\partial_t u^j - \eta^j X_F(t, u) - X_H(t, u^j)) &= 0 \\ \partial_s \eta^j - \int_0^1 F(t, u^j) dt &= 0 \end{aligned} \right\} \quad (5.6)$$

of energy

$$E(\varpi^j) \leq \limsup E(w_n) \leq \|H\|. \quad (5.7)$$

The sequence $w_n(s)$ converges to a solution $\varpi^0 = (u^0, \eta^0)$ of

$$\left. \begin{aligned} \partial_s u^0 + J(s, t, u^0) (\partial_t u^0 - \eta^0 X_F(t, u) - \beta_\infty^+(s) X_H(t, u^0)) &= 0 \\ \partial_s \eta^0 - \int_0^1 F(t, u^0) dt &= 0 \end{aligned} \right\} \quad (5.8)$$

and the sequence $w_n(s + (k+1)n)$ converges to a solution $\varpi^{k+1} = (u^{k+1}, \eta^{k+1})$ of

$$\left. \begin{aligned} \partial_s u^{k+1} + J(s, t, u^{k+1}) (\partial_t u^{k+1} - \eta^{k+1} X_F(t, u) - \beta_\infty^-(s) X_H(t, u^{k+1})) &= 0 \\ \partial_s \eta^{k+1} - \int_0^1 F(t, u^{k+1}) dt &= 0 \end{aligned} \right\} \quad (5.9)$$

where $\beta_\infty^\pm$ are defined at the beginning of section 3. In particular,

$$(y_\pm^j, \eta_\pm^j) := \varpi^j(\pm\infty) \in \text{Crit}\mathcal{A}^{(F, H)}, \quad j = 1, \dots, k \quad (5.10)$$

and

$$(y_+^0, \eta_+^0) := \varpi^0(+\infty), \quad (y_-^{k+1}, \eta_-^{k+1}) := \varpi^{k+1}(-\infty) \in \text{Crit}\mathcal{A}^{(F, H)} \quad (5.11)$$

are critical points. Thus, $y_{\pm}^j(0)$ are leaf-wise intersections, see Proposition 2.5. Moreover, they are ordered by action as follows

$$\begin{aligned} \mathcal{A}^{(F,H)}(y_+^0, \eta_+^0) &\geq \mathcal{A}^{(F,H)}(y_-^1, \eta_-^1) \geq \mathcal{A}^{(F,H)}(y_+^1, \eta_+^1) \geq \mathcal{A}^{(F,H)}(y_-^2, \eta_-^2) \geq \dots\dots \\ &\dots\dots \geq \mathcal{A}^{(F,H)}(y_+^k, \eta_+^k) \geq \mathcal{A}^{(F,H)}(y_-^{k+1}, \eta_-^{k+1}) . \end{aligned} \quad (5.12)$$

This follows directly from the choice of the sequences $w_n(s+jn)$. If $\mathcal{A}^{(F,H)}(\varpi_-^j) = \mathcal{A}^{(F,H)}(\varpi_+^j)$ then by Lemma 2.9 $E(\varpi^j) = \mathcal{A}^{(F,H)}(\varpi_-^j) - \mathcal{A}^{(F,H)}(\varpi_+^j) = 0$ and thus $\partial_s \varpi^j(s) = 0$. We recall that $\text{ev}_n(w_n) \in W^s(x_1, f_1) \times \dots \times W^u(x_k, f_k)$. By C_{loc}^∞ -convergence of $w_n(s+jn) \rightarrow \varpi^j(s) = (u^j, \eta^j)$ we conclude

$$u^j(0,0) \in \overline{W^s(x_j, f_j)} \quad (5.13)$$

where $\varpi^j = (u^j, \eta^j)$. Now, since $\partial_s \varpi^j(s) = 0$, i.e. $\varpi^j(s) = (y_-^j, \eta_-^j) = (y_+^j, \eta_+^j)$, we have

$$u^j(0,0) = y_-^j(0) = y_+^j(0) \in \overline{W^s(x_j, f_j)} . \quad (5.14)$$

But this contradicts assumption (1) that $W^s(x_j, f_j) \cap \mathcal{LI} = \emptyset$ since $\overline{\partial W^s(x_j, f_j)}$ is composed out of stable manifolds of critical points of higher indices. Therefore, $\partial_s \varpi^j \neq 0$ for all $j = 1, \dots, k$ and we conclude

$$\mathcal{A}^{(F,H)}(y_-^j, \eta_-^j) > \mathcal{A}^{(F,H)}(y_+^j, \eta_+^j), \quad j = 1, \dots, k . \quad (5.15)$$

If we denote $(y_l, \eta_l) := (y_-^l, \eta_-^l)$ for $l = 1, \dots, k$ and $(y_{k+1}, \eta_{k+1}) := (y_+^k, \eta_+^k)$ we have

$$\mathcal{A}^{(F,H)}(y_+^0, \eta_+^0) \geq \mathcal{A}^{(F,H)}(y_1, \eta_1) > \dots > \mathcal{A}^{(F,H)}(y_{k+1}, \eta_{k+1}) \geq \mathcal{A}^{(F,H)}(y_-^{k+1}, \eta_-^{k+1}) . \quad (5.16)$$

In particular, all (y_l, η_l) are different as critical points of $\mathcal{A}^{(F,H)}$. To finally conclude that the leaf-wise intersections $y_1(0), \dots, y_{k+1}(0)$ are all distinct we recall from Proposition 2.5 that two critical points of $\mathcal{A}^{(F,H)}$ can give rise to the same leaf-wise intersection only if some $y_j(0)$ is a periodic leaf-wise intersection. If $y_l(0) = y_{l'}(0)$ then as in the proof of [AF10a, Lemma 2.19] we conclude that

$$|\mathcal{A}^{(F,H)}(y_l, \eta_l) - \mathcal{A}^{(F,H)}(y_{l'}, \eta_{l'})| \geq \wp(\Sigma, \alpha) > \|H\| . \quad (5.17)$$

On the other hand we estimate for $(y_+^0, \eta_+^0) = \varpi^0(+\infty)$

$$\begin{aligned} \mathcal{A}^{(F,H)}(\varpi^0(+\infty)) - \mathcal{A}^{(F,H)}(\varpi^0(-\infty)) &= \int_{-\infty}^{\infty} \frac{d}{ds} \mathcal{A}^{(F, \beta_\infty^+(s)H)}(\varpi^0(s)) ds \\ &= \int_{-\infty}^{\infty} d\mathcal{A}^{(F,H)}(\varpi^0) \cdot \partial_s \varpi^0 ds + \int_{-\infty}^{\infty} \frac{\partial \mathcal{A}^{(F, \beta_\infty^+(s)H)}}{\partial s}(\varpi^0) ds \\ &= - \int_{-\infty}^{\infty} \|\partial_s \varpi^0\|^2 ds - \int_{-\infty}^{\infty} \int_0^1 \frac{\partial(\beta_\infty^+(s)H)}{\partial s}(t, u^0) dt ds \\ &\leq - \int_{-\infty}^{\infty} \underbrace{\left(\frac{\partial}{\partial s} \beta_\infty^+(s) \right)}_{\geq 0} \int_0^1 H(t, u^0(s, t)) dt ds \\ &\leq - \underbrace{\int_{-\infty}^{\infty} \left(\frac{\partial}{\partial s} \beta_\infty^+(s) \right) ds}_{=1} \cdot \int_0^1 \min_M H(t, \cdot) dt \\ &= - \int_0^1 \min_M H(t, \cdot) dt . \end{aligned}$$

The critical point $\varpi^0(-\infty)$ of $\mathcal{A}^{(F,0)}$ corresponds either to a Reeb orbit or constant loop in Σ . Since $E(\varpi^0) \leq \|H\|$ the same argument as in Proposition 3.2 rules out the former case. In particular, $\mathcal{A}^{(F,0)}(\varpi^0(-\infty)) = 0$ and therefore

$$\mathcal{A}^{(F,H)}(y_+^0, \eta_+^0) \leq - \int_0^1 \min_M H(t, \cdot) dt. \quad (5.18)$$

Analogously, for $(y_-^{k+1}, \eta_-^{k+1}) = \varpi^{k+1}(-\infty)$

$$\mathcal{A}^{(F,H)}(y_-^{k+1}, \eta_-^{k+1}) \geq - \int_0^1 \max_M H(t, \cdot) dt \quad (5.19)$$

is derived. Thus, if we assume $y_l(0) = y_{l'}(0)$ we have the following inequalities

$$\begin{aligned} \|H\| &= - \int_0^1 \min_M H(t, \cdot) dt + \int_0^1 \max_M H(t, \cdot) dt \\ &\geq \mathcal{A}^{(F,H)}(y_+^0, \eta_+^0) - \mathcal{A}^{(F,H)}(y_-^{k+1}, \eta_-^{k+1}) \\ &\geq |\mathcal{A}^{(F,H)}(y_l, \eta^l) - \mathcal{A}^{(F,H)}(y_{l'}, \eta^{l'})| \\ &\geq \wp(\Sigma, \alpha) \\ &> \|H\| \end{aligned} \quad (5.20)$$

where the second inequality follows from (5.16) and the last two from (5.17). This contradiction shows that the leaf-wise intersection points $y_1(0), \dots, y_{k+1}(0)$ are all distinct. This finishes the proof of Theorem 1.

Acknowledgments. This article was written during visits of the authors at the Institute for Advanced Study, Princeton. The authors thank the Institute for Advanced Study for their stimulating working atmospheres. The authors are grateful to Alberto Abbondandolo, Urs Frauenfelder, and Helmut Hofer for helpful discussions.

This material is based upon work supported by the National Science Foundation under agreement No. DMS-0635607 and DMS-0903856. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

REFERENCES

- [AF08] P. Albers and U. Frauenfelder, *Infinitely many leaf-wise intersections on cotangent bundles*, 2008, arXiv:0812.4426.
- [AF10a] ———, *Leaf-wise intersections and Rabinowitz Floer homology*, J. Topol. Anal. **2** (2010), no. 1, 77–98.
- [AF10b] ———, *On a Theorem by Ekeland-Hofer*, 2010, arXiv:1001.3386, to appear in Israel Journal of Mathematics.
- [AF10c] ———, *Spectral invariants in Rabinowitz Floer homology and global Hamiltonian perturbations*, 2010, arXiv:1001.2920.
- [AM09] P. Albers and M. McLean, *Non-displaceable contact embeddings and infinitely many leaf-wise intersections*, 2009, arXiv:0904.3564, to appear in Journal of Symplectic Geometry.
- [Ban80] A. Banyaga, *On fixed points of symplectic maps*, Invent. Math. **56** (1980), no. 3, 215–229.
- [CF09] K. Cieliebak and U. Frauenfelder, *A Floer homology for exact contact embeddings*, Pacific J. Math. **293** (2009), no. 2, 251–316.
- [Dra08] D. L. Dragnev, *Symplectic rigidity, symplectic fixed points, and global perturbations of Hamiltonian systems*, Comm. Pure Appl. Math. **61** (2008), no. 3, 346–370.
- [EH89] I. Ekeland and H. Hofer, *Two symplectic fixed-point theorems with applications to Hamiltonian dynamics*, J. Math. Pures Appl. (9) **68** (1989), no. 4, 467–489 (1990).

- [Flo89] A. Floer, *Cuplength estimates on Lagrangian intersections*, Comm. Pure Appl. Math. **42** (1989), no. 4, 335–356.
- [Gin07] V. L. Ginzburg, *Coisotropic intersections*, Duke Math. J. **140** (2007), no. 1, 111–163.
- [Gür09] B. Gürel, *Leafwise Coisotropic Intersections*, Int. Math. Res. Not. (2009), article ID rnp 164.
- [Hof88] H. Hofer, *Lusternik-Schnirelman-theory for Lagrangian intersections*, Ann. Inst. H. Poincaré Anal. Non Linéaire **5** (1988), no. 5, 465–499.
- [Hof90] ———, *On the topological properties of symplectic maps*, Proc. Roy. Soc. Edinburgh Sect. A **115** (1990), no. 1-2, 25–38.
- [Kan09] J. Kang, *Existence of leafwise intersection points in the unrestricted case*, 2009, arXiv:0910.2369.
- [Liu05] C.-G. Liu, *Cup-length estimate for Lagrangian intersections*, J. Differential Equations **209** (2005), no. 1, 57–76.
- [Mer10] W. Merry, *On the Rabinowitz Floer homology of twisted cotangent bundles*, 2010, arXiv:1002.0162.
- [Mos78] J. Moser, *A fixed point theorem in symplectic geometry*, Acta Math. **141** (1978), no. 1–2, 17–34.
- [MS98] D. McDuff and D. A. Salamon, *Introduction to symplectic topology*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1998.
- [Sch93] M. Schwarz, *Morse homology*, Progress in Mathematics, vol. 111, Birkhäuser Verlag, Basel, 1993.
- [Sch98] ———, *A quantum cup-length estimate for symplectic fixed points*, Invent. Math. **133** (1998), no. 2, 353–397.
- [Zil08] F. Ziltener, *Coisotropic Submanifolds, Leafwise Fixed Points, and Presymplectic Embeddings*, 2008, arXiv:0811.3715, to appear in Journal of Symplectic Geometry.

PETER ALBERS, DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY
E-mail address: `palbers@math.purdue.edu`

AL MOMIN, DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY
E-mail address: `amomin@math.purdue.edu`